

2

AD-A209 277

OFFICE OF NAVAL RESEARCH

Contract N00014-87-K-0541

R & T Code 4133020

Technical Report No. 13

The Adsorption of Isobutyramide at the Mercury/Aqueous Sodium Fluoride Solution Interface

by

W.R. Fawcett and A.J. Motheo

Prepared for Presentation

at

Electrochemical Society Meeting  
Los Angeles, CA May 1989


May 1, 1989

DTIC  
ELECTE  
JUN 20 1989  
S E D

Reproduction in whole or in part is permitted  
for any purpose of the United States Government

"This document has been approved for public release  
and sale; its distribution is unlimited"

89 6 10 0 12

<b>REPORT DOCUMENTATION PAGE</b>		<b>1. REPORT NO.</b> 13	<b>2.</b>	<b>3. Revision's Accession No.</b>
<b>4. Title and Subtitle</b> The Adsorption of Isobutyramide at the Mercury/Aqueous Sodium Fluoride Solution Interface				<b>5. Report Date</b> May 1, 1989
<b>7. Author(s)</b> W.R. Fawcett and A.J. Motheo				<b>6.</b>
<b>8. Performing Organization Name and Address</b> Department of Chemistry University of California Davis, CA 95616				<b>9. Performing Organization Report No.</b>
<b>10. Project/Task/Work Unit No.</b>				<b>11. Contract/Grant No.</b> (a) N00014-87-K-0541 (b)
<b>12. Sponsoring Organization Name and Address</b> Office of Naval Research 800 N. Quincy Arlington, VA 22217-5000				<b>13. Type of Report &amp; Period Covered</b>
<b>15. Supplementary Notes</b> Prepared for presentation at: Electrochemical Society Meeting, Los Angeles, CA, May 1989.				<b>14.</b>
<b>16. Abstract (Limit 200 words)</b> The differential capacity of the electrode/solution interface was measured at constant concentration of NaF (0.25 m) for iso-butyramide concentrations in the range 0.01 to 1 m. The capacity curves were typical of those with moderately strong organic adsorption with a deep shallow minimum negative of the p.z.c. and desorption peaks at extreme positive and negative potentials. Analysis of these data demonstrates that reorientation of the adsorbed amide molecule plays an important role in determining the characteristics of the adsorption isotherm. (JES) 				
<b>17. Document Analysis a. Descriptors</b>				
<b>b. Identifiers/Open-Ended Terms</b>				
<b>c. CDBAT Field/Group</b>				
<b>18. Availability Statement</b>		<b>19. Security Class (This Report)</b> Unclassified		<b>21. No. of Pages</b>
		<b>20. Security Class (This Page)</b>		<b>22. Price</b>

# 75-Word Abstract Form

Extended Abstract must be submitted with the 75-Word Abstract by December 1, 1988

**Los Angeles, California—May 7-12, 1989**

Submit to: The Electrochemical Society, Inc.  
10 South Main Street, Pennington, NJ 08534-2896  
With a copy to the Organizing Chairman

Abstract No. ....  
(to be assigned by the Society)

Schedule for ..... General Session--  
(Title of Symposium)

Sponsored by ..... Physical Electrochemistry  
(Division/Group)

Title of paper ..... The Adsorption of Isobutyramide at the Mercury/Aqueous Sodium Fluoride  
Solution Interface

Authors (Underline name of author presenting paper.) ..... W. Ronald Fawcett and Artur J. Motheo

Business affiliation and address ..... Department of Chemistry  
University of California  
Davis, CA 95616 (916) 752-1105  
(State or Country) (ZIP Code) (Telephone No.)

(Type abstract in this area—double-spaced.)

The differential capacity of the electrode/solution interface was measured at constant concentration of NaF (0.25 m) for iso-butyramide concentrations in the range 0.01 to 1 m. The capacity curves were typical of those with moderately strong organic adsorption with a deep shallow minimum negative of the p.z.c. and desorption peaks at extreme positive and negative potentials. Analysis of these data demonstrates that reorientation of the adsorbed amide molecule plays an important role in determining the characteristics of the adsorption isotherm.

Do you require any audiovisual equipment?

- ☐ 35 mm (2 x 2 in.) slide projector  
☒ Overhead projector  
☐ Specify other (subject to availability and cost)

Has the information in this abstract been presented verbally, submitted for publication, or published?

☐ Yes ☐ No

If the answer is yes, please provide the reference (except in the case of invited review presentations).

Is a full length paper on this work to be submitted for Society Journal publication? ☐ Yes ☐ No

Papers presented before a Society technical meeting become the property of the Society and may not be published elsewhere without written permission of the Society. Papers presented at Society technical meetings must be authored by a member or sponsored by an active member.

W.R. Fawcett, A.J. Motheo

The Adsorption of Isobutyramide at the Mercury/Aqueous Sodium Fluoride Solution Interface.

W. Ronald Fawcett and Amur J. Motheo\*

Department of Chemistry, University of California, Davis, CA 95616

We have recently been involved in a study of the adsorption of amides from aqueous solution at the mercury electrode [1]. The adsorptive behavior of these molecules is quite different from other organic systems, displaying characteristics which can be attributed to molecular reorientation. Some of these unusual interfacial properties can be attributed to the fact that these molecules have quite large dipole moments. The present study was concerned with the adsorption of isobutyramide (IBA) which can be classified as moderately strong compared to the adsorption of acetamide and dimethylacetamide [1].

The adsorption of IBA was studied by measuring the differential capacity of the mercury/solution interface as a function of electrode potential and adsorbate concentration, holding the concentration of the supporting electrolyte constant (0.25 M NaF). The reference electrode was a saturated calomel electrode (SCE), and the potential of zero charge was obtained using the streaming mercury electrode technique. The change in salt activity with IBA concentration was monitored with specific ion electrodes. These measurements also provided the factors to convert the SCE potential scale to one of thermodynamic significance. All experiments were conducted at 25°C.

Differential capacity against potential data for IBA concentrations in the range 0.01 to 1.0 M are shown in Fig. 1. These curves display a minimum at potentials negative of the p.z.c., whose depth increases with IBA concentration. The maximum at more negative potentials is attributed to molecular desorption. The p.z.c. was observed to move in the positive direction with increase in IBA concentration, an observation which is attributed to adsorption of the molecular dipole with its positive end towards the metal in this potential region.

The capacity curves were twice integrated with respect to potential to obtain the relative surface tension. These data were then used to obtain the relative surface excess of IBA,  $\Gamma$  as a function of bulk activity for constant electrode charge density. The results, shown in Fig. 2, indicate that  $\Gamma$  reaches a maximum value at  $-4 \mu\text{C cm}^{-2}$  for high bulk concentrations; however, at lower concentrations,  $\Gamma$  slowly increases as the electrode charge density is made more negative. Parry and Parsons [2] found similar behavior in adsorption of the p-toluene sulphonate anion at mercury and proposed a two-position isotherm to describe surface coverage. An analysis of the present data shows that reorientation of the IBA molecule occurs in the potential range in

which adsorption is significant. The results of this analysis and the parameters of the adsorption isotherm are discussed in detail in this paper. The discussion also includes an analysis of the effects of ignoring variation in salt activity with increase in IBA concentration.

References

1. W.R. Fawcett, G.Y. Champagne, S. Kono and A.J. Motheo, *J. Phys. Chem.*, **92**, 6368 (1988).
2. J.M. Parry and R. Parsons, *J. Electrochem. Soc.*, **113**, 992 (1966).

\*Permanent address - Instituto de Física e Química de São Carlos, USP, São Carlos, SP 13560, Brazil

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	




Fig. 1. Differential capacity against electrode potential for the Hg/solution interface for the system 0.25 m NaF +  $x$  m isobutyramide where  $x$  is equal to: 0(1), 0.01(2), 0.018(3), 0.032(4), 0.056(5), 0.10(6), 0.18(7), 0.32(8), 0.56(9) and 1.00(10).




Fig. 2. Surface excess of isobutyramide against electrode charge density for various bulk concentrations: 0.01(1), 0.032(2), 0.10(3), 0.18(4), 0.32(5) and 0.56 m (6).